

Exact Four-Dimensional Dyonic Black Holes and Bertotti-Robinson Spacetimes in String Theory

David A. Lowe*

and

Andrew Strominger†

Department of Physics
University of California
Santa Barbara, CA 93106-9530

Conformal field theories corresponding to two-dimensional electrically charged black holes and to two-dimensional anti-de Sitter space with a covariantly constant electric field are simply constructed as $SL(2, \mathbb{R})/\mathbb{Z}$ WZW coset models. The two-dimensional electrically charged black holes are related by Kaluza–Klein reduction to the 2+1-dimensional rotating black hole of Bañados, Teitelboim and Zanelli, and our construction is correspondingly related to its realization as a WZW model. Four-dimensional spacetime solutions are obtained by tensoring these two-dimensional theories with $SU(2)/Z(m)$ coset models. These describe a family of dyonic black holes and the Bertotti–Robinson universe.

March, 1994

* lowe@tpau.physics.ucsb.edu

† andy@denali.physics.ucsb.edu

1. Introduction

Near extremality, four-dimensional, magnetically charged black holes in string theory develop a large cigar-shaped region adjacent to the horizon [1]. The solution in this region degenerates into a product of two two-dimensional factors, both of which can be described as exact conformal field theories and hence as exact solutions of string theory. One of these factors is Witten's two-dimensional black hole [2], and the other is a two-sphere with magnetic charge and is represented by a WZW orbifold [3,4]. At extremality, the two-dimensional black hole factor is replaced by the linear dilaton vacuum.

It is known from analysis of the low-energy effective field theory that the addition of electric charge dramatically changes the picture [1,5,6]. The solutions still degenerate into two factors near the horizon, but the two-dimensional black hole factor of [2] must be replaced by an asymptotically anti-de Sitter, electrically charged black hole similar to those discussed in [7]. In the extremal limit, the black hole factor becomes electrically charged anti-de Sitter space. Including the magnetically charged two-sphere, one then has the Bertotti-Robinson universe. This limiting behavior is identical to that of Reissner-Nordström black holes.

In this paper, we give an exact construction of the electrically charged two-dimensional black hole, and corresponding family of four-dimensional dyonic black holes. This is rather easily accomplished by using the observation of Achúcarro and Ortiz [8] that the 2+1-dimensional rotating black hole of Bañados, Teitelboim and Zanelli [9] may be reinterpreted as an electrically charged two-dimensional black hole. We may then apply the exact construction of those black holes by Horowitz, Welch and Kaloper [10] to our case. Related constructions give two-dimensional electrically charged anti-de Sitter space and the Bertotti-Robinson universe.

2. Kaluza-Klein reduction from 3 to 2 dimensions

2.1. Two-dimensional charged black holes

Bañados et al. [9] have constructed a family of spinning black holes in 2 + 1-dimensional Einstein gravity with negative cosmological constant proportional to l^2 . For general values of the mass M and angular momentum J , the 2 + 1-dimensional spacetime metric is

$$ds^2 = \left(M - \frac{r^2}{l^2}\right) dt^2 - J dt d\varphi + r^2 d\varphi^2 + \left(\frac{r^2}{l^2} - M + \frac{J^2}{4r^2}\right)^{-1} dr^2, \quad (2.1)$$

where φ is periodically identified with period 2π , and r and t may take any real value. There are two horizons located at $r = r_{\pm}$, where $r_{\pm}$ are related to M and J via

$$M = \frac{r_+^2 + r_-^2}{l^2}, \quad J = \frac{2r_+ r_-}{l}. \quad (2.2)$$

The above identification breaks down in the extreme case $|J| = Ml$, when the two horizons coincide, and also in the massless case $M = J = 0$. In these situations, the periodic identification must be made along a null direction.

The 2+1 black hole (2.1) also corresponds to a solution of string theory with a non-trivial antisymmetric tensor [10]. The relevant part of the three-dimensional string effective action is

$$S_3 = \int d^3x \sqrt{-\hat{g}} e^{-2\hat{\phi}} \left[\frac{4}{l^2} + \hat{R} + 4(\nabla\hat{\phi})^2 - \frac{1}{12}\hat{H}^2 \right], \quad (2.3)$$

where $3l^2/(l^2 - 2)$ is the central charge of the conformal field theory. The antisymmetric tensor field strength for the black hole solutions is

$$\hat{H}_{\hat{\mu}\hat{\nu}\hat{\lambda}} = \frac{2}{l} \epsilon_{\hat{\mu}\hat{\nu}\hat{\lambda}}, \quad (2.4)$$

where $\hat{\mu}$ runs over r, t, φ .

Here we would like to reinterpret this three-dimensional solution as a two-dimensional charged black hole. This is achieved, following Achúcarro and Ortiz, by performing a Kaluza–Klein reduction of the metric (2.1) and antisymmetric tensor, regarding φ as the compact coordinate which will generate a $U(1)$ gauge field. The 2 + 1-dimensional space-time metric is written in terms of the two-dimensional metric $g_{\mu\nu}$, $U(1)$ gauge field A_μ , and scalar D

$$\hat{g} = \begin{pmatrix} g_{\mu\nu} + e^{2D} A_\mu A_\nu & e^{2D} A_\mu \\ e^{2D} A_\nu & e^{2D} \end{pmatrix}. \quad (2.5)$$

Here μ, ν run over r, t and unhatted quantities refer to two-dimensional fields. The two-dimensional low-energy effective action is

$$S_2 = \int d^2x \sqrt{-g} e^{D-2\hat{\phi}} \left[\frac{4}{l^2} + R + 4(\nabla\hat{\phi})^2 - 4\nabla D \cdot \nabla\hat{\phi} - \frac{1}{4}e^{2D}F^2 - \frac{1}{4}e^{-2D}F'^2 \right], \quad (2.6)$$

where F (F') is the field strength constructed from A_μ ($\hat{B}_{\mu\varphi}$). The antisymmetric tensor $B_{\mu\nu}$ drops out of this expression and will henceforth be ignored.

It is possible to eliminate one of the gauge fields and the dilaton in a manner consistent with the equations of motion. This is achieved by setting

$$\hat{\phi} = 0, \quad F'_{\mu\nu} = \frac{2e^D}{l} \epsilon_{\mu\nu}, \quad (2.7)$$

where $\epsilon_{\mu\nu}$ is the two-dimensional volume form. The truncated equations of motion then follow from

$$S_2 = \int d^2x \sqrt{-g} e^D \left[\frac{2}{l^2} + R - \frac{1}{4}e^{2D}F^2 \right]. \quad (2.8)$$

The two-dimensional spacetime solutions we consider are all compatible with this truncation. (2.8) is the Jackiw–Teitelboim model with a gauge field [11].

The Kaluza–Klein reduction of the three-dimensional metric (2.1) and antisymmetric tensor field leads to

$$ds^2 = \left(M - \frac{r^2}{l^2} - \frac{J^2}{4r^2} \right) dt^2 + \left(\frac{r^2}{l^2} - M + \frac{J^2}{4r^2} \right)^{-1} dr^2, \quad (2.9)$$

with $U(1)$ gauge field and scalar

$$A_t = -\frac{J}{2r^2} , \quad D = \log r , \quad (2.10)$$

which is of course an extrema of (2.8).¹

Evaluating the Ricci scalar for the metric (2.9) gives

$$R = -\frac{4r^4 + 3J^2l^2}{2l^2r^4} . \quad (2.11)$$

The two-dimensional black hole metric is asymptotically anti-de Sitter, with a curvature singularity at $r = 0$.

Horowitz, Welch and Kaloper [10] have shown that the solution (2.4), (2.1), corresponds to the $SL(2, \mathbb{R})$ WZW model with a discrete identification and is therefore exact. The $SL(2, \mathbb{R})$ WZW action is invariant under the following symmetries

$$\delta g = \epsilon \left[\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} g + g \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right] \quad (2.12)$$

and

$$\delta g = \epsilon \left[\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} g - g \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \right] , \quad (2.13)$$

where g is an element of $SL(2, \mathbb{R})$. Quotienting $SL(2, \mathbb{R})$ by a discrete subgroup of a linear combination of these symmetries yields the general two parameter family of black holes (2.1). The level k_{SL} of the WZW model is fixed by

$$k_{SL} = l^2 . \quad (2.14)$$

2.2. 1+1-dimensional anti-de Sitter space with constant electric field

Anti-de Sitter space in 1+1 dimensions with a covariantly constant electric field (not considered in [10]) may be obtained by a similar construction which we will describe in some detail. The difference here will be that the periodic identification of 2+1-dimensional anti-de Sitter space (*ie* $SL(2, \mathbb{R})$) will be made along a direction that is always spacelike. In the previous section the identification was made along a Killing vector that was timelike in certain regions.

¹ Note that decompactification ($D \rightarrow \infty$) occurs at large r . However, in order to obtain an asymptotically flat four-dimensional black hole one must tensor with a two-sphere as discussed in subsequent sections and “sew on” an asymptotically flat four-dimensional region at some finite radius from the horizon. The growth of D will then be cut off at the sewing radius. This sewing procedure is readily understood in the low-energy effective field theory [6,12], but not in the conformal field theory, where it presumably corresponds to perturbation by a “bad” marginal operator.

It is convenient to use coordinates which differ from those of the previous subsection to describe the anti-de Sitter solution. Here we parametrize an element of $SL(2, \mathbb{R})$, with all elements non-zero, by [13]

$$g = d_1(-e)^{\epsilon_1} s^{\epsilon_2} p d_2 , \quad (2.15)$$

where $\epsilon_1, \epsilon_2 = 0$ or 1 ,

$$\begin{aligned} d_1 &= \begin{pmatrix} e^{t_L/2} & 0 \\ 0 & e^{-t_L/2} \end{pmatrix} , & d_2 &= \begin{pmatrix} e^{(-1)^{\epsilon_2} t_R/2} & 0 \\ 0 & e^{-(-1)^{\epsilon_2} t_R/2} \end{pmatrix} , \\ -e &= \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} & s &= \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} , \end{aligned} \quad (2.16)$$

and p is either

$$\begin{aligned} (i) \quad p &= \begin{pmatrix} \cosh(r/2) & \sinh(r/2) \\ \sinh(r/2) & \cosh(r/2) \end{pmatrix} , & -\infty < r < \infty & \text{ or} \\ (ii) \quad p &= \begin{pmatrix} \cos(r/2) & \sin(r/2) \\ -\sin(r/2) & \cos(r/2) \end{pmatrix} , & -\pi/2 < r < \pi/2 , \end{aligned} \quad (2.17)$$

with $t_L, t_R \in \mathbb{R}$.

The WZW action is

$$S_{WZW} = \frac{k_{SL}}{4\pi} \int_{\Sigma} d^2 z \text{Tr}(g^{-1} \partial g g^{-1} \bar{\partial} g) + k_{SL} \Gamma(g) , \quad (2.18)$$

where Γ is the Wess Zumino term

$$\Gamma = \frac{1}{12\pi} \int_B \text{Tr}(g^{-1} dg \wedge g^{-1} dg \wedge g^{-1} dg) . \quad (2.19)$$

Here Σ is a Riemann surface and B is a 3-manifold with boundary Σ . With p in region (i), (2.18) takes the form

$$S_{WZW} = \frac{k_{SL}}{4\pi} \int (\partial r \bar{\partial} r + \partial t_L \bar{\partial} t_L + \partial t_R \bar{\partial} t_R + 2 \cosh r \bar{\partial} t_L \partial t_R) . \quad (2.20)$$

With p in region (ii) the answer is

$$S_{WZW} = \frac{k_{SL}}{4\pi} \int (-\partial r \bar{\partial} r + \partial t_L \bar{\partial} t_L + \partial t_R \bar{\partial} t_R + 2 \cos r \bar{\partial} t_L \partial t_R) . \quad (2.21)$$

This may be viewed as a σ -model on a three-dimensional target space with metric $\hat{g}_{\hat{\mu}\hat{\nu}}$ and antisymmetric tensor $\hat{B}_{\hat{\mu}\hat{\nu}}$. One finds in region (i)

$$\hat{g} = \frac{1}{4} \begin{pmatrix} k_{SL} & 0 & 0 \\ 0 & k_{SL} & k_{SL} \cosh r \\ 0 & k_{SL} \cosh r & k_{SL} \end{pmatrix} , \quad (2.22)$$

with the ordering of coordinates taken to be (r, t_L, t_R) . The non-zero component of the antisymmetric tensor is

$$\hat{B}_{t_L t_R} = \frac{1}{4} k_{SL} \cosh r . \quad (2.23)$$

In region (ii) one finds

$$\hat{g} = \frac{1}{4} \begin{pmatrix} -k_{SL} & 0 & 0 \\ 0 & k_{SL} & k_{SL} \cos r \\ 0 & k_{SL} \cos r & k_{SL} \end{pmatrix} , \quad (2.24)$$

with antisymmetric tensor

$$\hat{B}_{t_L t_R} = \frac{1}{4} k_{SL} \cos r . \quad (2.25)$$

The WZW action is invariant under translations in t_R which are generated by

$$\delta g = \epsilon g \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} . \quad (2.26)$$

We quotient $SL(2, \mathbb{R})$ by a discrete subgroup of this symmetry: the integers $\mathbb{Z}$. t_R becomes a periodic coordinate with range $[0, 2\pi/\gamma)$, where γ is a real parameter.

We may now perform a Kaluza–Klein reduction of the metric and antisymmetric tensor, to obtain a two–dimensional spacetime with a $U(1)$ gauge field in the same way as before. The identifications of (2.5) give the fields in region (i)

$$\begin{aligned} A_{t_L} &= \gamma \cosh r , & e^{2D} &= \frac{k_{SL}}{4\gamma^2} , \\ g &= \frac{1}{4} \begin{pmatrix} k_{SL} & 0 \\ 0 & -k_{SL} \sinh^2 r \end{pmatrix} . \end{aligned} \quad (2.27)$$

Similarly in region (ii) the fields are

$$\begin{aligned} A_{t_L} &= \gamma \cos r , & e^{2D} &= \frac{k_{SL}}{4\gamma^2} , \\ g &= \frac{1}{4} \begin{pmatrix} -k_{SL} & 0 \\ 0 & k_{SL} \sin^2 r \end{pmatrix} . \end{aligned} \quad (2.28)$$

We now show this corresponds to a double cover of two–dimensional anti–de Sitter space, $(adS)_2$. Recall $(adS)_2$ may be defined as the hyperboloid

$$-x_0^2 - x_1^2 + x_2^2 = -1 , \quad (2.29)$$

embedded in flat three–dimensional space with metric

$$ds^2 = -dx_0^2 - dx_1^2 + dx_2^2 . \quad (2.30)$$

Now define coordinates

$$\begin{aligned}x_0 &= \cosh r \\x_1 &= \sinh r \sinh t_L \\x_2 &= \sinh r \cosh t_L ,\end{aligned}\tag{2.31}$$

which covers the region $x_0 > 1$ when $r, t_L \in \mathbb{R}$. The metric in these coordinates is

$$ds^2 = dr^2 - \sinh^2 r dt_L^2 ,\tag{2.32}$$

as in region (i), up to a rescaling by $k_{SL}/4$. Remember we actually have 4 copies of region (i) corresponding to $\epsilon_1 = 0, 1$ and $\epsilon_2 = 0, 1$. Two of these copies are sufficient to cover the region $|x_0| > 1$.

To cover the region $0 < x_0 < 1$ define coordinates

$$\begin{aligned}x_0 &= \cos r \\x_1 &= \sin r \cosh t_L \\x_2 &= \sin r \sinh t_L ,\end{aligned}\tag{2.33}$$

where $t_L \in \mathbb{R}$ and $r \in (-\pi/2, \pi/2)$. The metric is then

$$ds^2 = -dr^2 + \sin^2 r dt_L^2 ,\tag{2.34}$$

agreeing with the metric above in region (ii). A second copy of region (ii) will cover $-1 < x_0 < 0$. We find then that the WZW model corresponds to a double cover of $(adS)_2$. It may also be verified that the gauge field strength is covariantly constant with respect to this metric.

3. Four-dimensional exact solutions

Having obtained the above two-dimensional spacetime solutions it is a simple matter to tensor these with the angular magnetic monopole CFT of [3,4] to obtain four-dimensional spacetime solutions with non-trivial electric and magnetic fields. The monopole CFT is obtained by quotienting a $SU(2)$ WZW model by the discrete subgroup $Z(m)$.

To parametrize the $SU(2)$ group manifold we introduce the real coordinates θ, ϕ and ζ and write the group element g' as

$$g' = e^{\frac{i}{2}\phi\sigma_3} e^{\frac{i}{2}\theta\sigma_1} e^{\frac{i}{2}\zeta\sigma_3} ,\tag{3.1}$$

where $\theta \in [0, \pi]$, $\phi \in [0, 2\pi]$ and $\zeta \in [0, 4\pi]$, and the σ_i are the Pauli matrices. The $SU(2)$ WZW action is then

$$S_{WZW} = \frac{k_{SU}}{4\pi} \int (\partial\theta\bar{\partial}\theta + \partial\phi\bar{\partial}\phi + \partial\zeta\bar{\partial}\zeta + 2\cos\theta \bar{\partial}\phi\partial\zeta) .\tag{3.2}$$

As in [4], this three-dimensional sigma model action may be compactified to 2 dimensions. The extra coordinate ζ , leads to a $U(1)$ gauge field corresponding to a magnetic field, while the two-dimensional space is just S^2 . Quotienting the $SU(2)$ manifold by a discrete subgroup $Z(m)$, identifying $\zeta \sim \zeta + \frac{4\pi}{m}$, determines the charge of the magnetic monopole $Q = m$. Modular invariance requires $k_{SU} = nm$ where n and m are integers. (Alternatively one could use the conformal field theories of [3], which involve magnetic charge associated with a general $U(1)$ current algebra.)

3.1. Four-dimensional charged black holes

Tensoring this monopole CFT with the CFT described in section 2.1 leads to a solution describing the throat limit of a four-dimensional black hole with electric and magnetic charge. The four-dimensional spacetime is identified with the r, t, θ and ϕ coordinates. The two compactified dimensions correspond to the ζ and φ coordinates.

In the following A_μ^φ (A_μ^ζ) refers to the gauge field derived from the $\hat{g}_{\mu\varphi}$ ($\hat{g}_{\mu\zeta}$) components of the six-dimensional metric. Also, $A'_{\mu,\varphi}$ ($A'_{\mu,\zeta}$) refers to the gauge field constructed from the $\hat{B}_{\mu\varphi}$ ($\hat{B}_{\mu\zeta}$) components of the six-dimensional antisymmetric tensor. The scalar field D_φ (D_ζ) is derived from the $\hat{g}_{\varphi\varphi}$ ($\hat{g}_{\zeta\zeta}$) component of the metric.

Note that in four-dimensions we may no longer truncate the gauge field $A'_{\mu,\varphi}$ and dilaton as in (2.7). However the gauge field $A'_{\mu,\zeta}$ and the scalar D_ζ may be truncated with the identification

$$D_\zeta = \text{const}, \quad A'_{\mu,\zeta} = \pm e^{2D_\zeta} A_\mu^\zeta, \quad (3.3)$$

in a manner consistent with the equations of motion. The non-trivial components of the spacetime fields are then

$$\begin{aligned} ds^2 &= \left(M - \frac{r^2}{l^2} - \frac{J^2}{4r^2} \right) dt^2 + \left(\frac{r^2}{l^2} - M + \frac{J^2}{4r^2} \right)^{-1} dr^2 + \frac{1}{4} k_{SU} d\theta^2 + \frac{1}{4} k_{SU} \sin^2 \theta d\phi^2, \\ A_t^\varphi &= -\frac{J}{2r^2}, \quad A'_{t,\varphi} = -\frac{r^2}{l}, \quad A_\phi^\zeta = m \cos \theta, \quad \hat{\phi} = \text{const}, \\ e^{2D_\varphi} &= r^2, \quad e^{2D_\zeta} = \frac{n}{4m}, \quad H_{\mu\nu\rho} = 0. \end{aligned} \quad (3.4)$$

Since we wish to identify this with a six-dimensional Kaluza-Klein spacetime, the central charge should be 6, i.e.

$$c = \frac{3k_{SU}}{k_{SU} + 2} + \frac{3k_{SL}}{k_{SL} - 2} = 6. \quad (3.5)$$

This leads to the condition $k_{SL} = k_{SU} + 4$. Taking k_{SL} and k_{SU} both positive leads to a spacetime with the correct $(-+++)$ signature.

3.2. Bertotti-Robinson Universe

If instead we tensor the monopole CFT with the anti-de Sitter solution of section 2.2, we will obtain, in the large k_{SU} and large k_{SL} limit, the four-dimensional Bertotti-

Robinson spacetime [14] with covariantly constant electric and magnetic fields. This solution describes the throat limit of the extremal dilaton black holes with electric and magnetic charge described in [6], and also the throat limit of the familiar extremal Reissner–Nordström black hole.

Here four-dimensional spacetime corresponds to the coordinates r, t_L, θ and ϕ while the two compactified dimensions correspond to t_R and ζ . The additional gauge field $A'_{\mu, \zeta}$ may be truncated as in (3.3), and A'_{μ, t_R} may be truncated in an analogous way by setting

$$D_{t_R} = \text{const}, \quad A'_{\mu, t_R} = \pm e^{2D_{t_R}} A_{\mu}^{t_R}. \quad (3.6)$$

The four-dimensional spacetime fields are then (for simplicity we write these only for region (i) of $SL(2, \mathbb{R})$)

$$\begin{aligned} ds^2 &= \frac{1}{4} k_{SL} dr^2 - \frac{1}{4} k_{SL} \sinh^2 r dt_L^2 + \frac{1}{4} k_{SU} d\theta^2 + \frac{1}{4} k_{SU} \sin^2 \theta d\phi^2, \\ A_{t_L}^{t_R} &= \gamma \cosh r, \quad A_{\phi}^{\zeta} = m \cos \theta, \quad \hat{\phi} = \text{const} \\ e^{2D_{t_R}} &= \frac{k_{SL}}{4\gamma^2}, \quad e^{2D_{\zeta}} = \frac{n}{4m}, \quad H_{\mu\nu\rho} = 0. \end{aligned} \quad (3.7)$$

The central charge should again be set to 6, which leads to $k_{SL} = k_{SU} + 4$.

As described in [6] this solution possesses $N = 2$ supersymmetry when embedded in $N = 4, d = 4$ supergravity which in turn is the low energy effective theory of the heterotic string with the $E_8 \times E_8$ gauge fields truncated. This corresponds to $(4, 0)$ worldsheet supersymmetry. In terms of the CFT, this is achieved by supersymmetrizing the WZW models by adding free fermions. The $(4, 0)$ supersymmetry algebra is then constructed as in [15,16].

Acknowledgements:

D. L. was supported in part by NSF grant PHY-91-16964, and A. S. was supported in part by DOE grant 91-ER40618.

References

- [1] G.W. Gibbons, *Nucl. Phys.* **B207** (1982) 337; G.W. Gibbons and K. Maeda, *Nucl. Phys.* **B298** (1988) 741; D. Garfinkle, G. Horowitz, and A. Strominger, *Phys. Rev.* **D43** (1991) 3140.
- [2] E. Witten, *Phys. Rev.* **D44** (1991) 314.
- [3] S. Giddings, J. Polchinski and A. Strominger, *Phys. Rev.* **D48** (1993) 5784.
- [4] W. Nelson, University of California at Santa Barbara preprint, UCSBTH-93-10.
- [5] A. Shapere, S. Trivedi and F. Wilczek, *Mod. Phys. Lett.* **A6** (1991) 337.
- [6] R. Kallosh, A. Linde, T. Ortin, A. Peet and A. Van Proeyen, *Phys. Rev.* **D46** (1992) 5278; R. Kallosh and A. Peet, *Phys. Rev.* **D46** (1992) 5223.
- [7] C. Nappi, M.D. McGuigan and S. Yost, *Nucl. Phys.* **B375** (1992) 421.
- [8] A. Achúcarro and M. Ortiz, *Phys. Rev.* **D48** (1993) 3600.
- [9] M. Bañados, C. Teitelboim, J. Zanelli, *Phys. Rev. Lett.* **69** (1992) 1849 ; M. Bañados, M. Henneaux, C. Teitelboim, J. Zanelli, *Phys. Rev.* **D 48** (1993) 1506.
- [10] G. Horowitz and D. Welch, *Phys. Rev. Lett.* **71** (1993) 328; N. Kaloper, *Phys. Rev.* **D48** (1993) 2598.
- [11] R. Jackiw, in *Quantum Theory of Gravity*, ed. S. Christensen, Hilger, Bristol 1984; C. Teitelboim, *ibid.*
- [12] S. Giddings and A. Strominger, *Phys. Rev.* **D46** (1992) 627.
- [13] N. Vilenkin and A. Klimyk, “Representation of Lie Groups and Special Functions,” Kluwer Academic Publishers (1991).
- [14] B. Bertotti, *Phys. Rev.* **116** (1959) 1331; I. Robinson, *Bull. Acad. Polon. Sci.* **7** (1959) 351.
- [15] A. Sevrin, W. Troost, and A. Van Proeyen, *Phys. Lett.* **208B** (1988) 447.
- [16] C.G. Callan, J.A. Harvey and A. Strominger, *Nucl. Phys.* **B359** (1991) 611.